
Cell Growthand Division

Cells have their own life cycle that includes reproduction, growth, and regulation, which allows organisms to carry out life functions and grow.

- 5.1 The Cell Cycle TEKS 5A
- 5.2 Mitosis and Cytokinesis TEKS 5A

Data Analysis
CONSTRUCTING DATA TABLES TEKS 2G

- 5.3 Regulation of the Cell Cycle
 TEKS 5A, 5B, 5C, 5D, 9C
 - **5.4** Asexual Reproduction
- 5.5 Multicellular Life TEKS 5B, 5C, 10C



ONLINE BIOLOGY

HMDScience.com

ONLINE Labs

- Mitosis in Onion Root Cells
- Quick Lab Cancer
- Modeling Cell Surface Area—to-Volume Ratio
- Apoptosis
- Animating Mitosis
- Virtual Lab Investigating Bacterial Growth

- Video Lab Mitosis in Plant Cells
- S.T.E.M. Lab Modeling Induction in Embryos



USING LANGUAGE

Cause and Effect In biological processes, one step leads to another step. When reading, you can often recognize these cause-and-effect relationships by words that indicate a result, such as so, consequently, if-then, and as a result.

YOUR TURN

Identify the cause and effect in the following sentences.

- 1. People often shiver as a result of being cold.
- 2. The light got brighter, so the pupil of the eye got smaller.
- **3.** If the cell passes the G_2 checkpoint, then the cell may begin to divide.



The Cell Cycle



VOCABULARY

cell cycle mitosis cytokinesis

TEKS 5A describe the stages of the cell cycle, including deoxyribonucleic acid (DNA) replication and mitosis, and the importance of the cell cycle to the growth of organisms

KEY CONCEPT Cells have distinct phases of growth, reproduction, and normal functions.

MAIN IDEAS

- The cell cycle has four main stages.
- Cells divide at different rates.
- O Cell size is limited.

Connect to Your World

Many of life's little chores can be quietly satisfying and rather fun. Washing dishes by hand, however, is not so fun, which is why some clever person made the dishwasher. This handy invention soaks, washes, and rinses your dishes to a spot-free, sanitary sparkle. You unload the dishes, and the machine is ready to start the cycle all over again. A cell goes through a cycle, too. This cycle of growth, DNA synthesis, and division is essential for an organism to grow and heal. If it goes out of control, abnormal cell growth may occur, resulting in cancer cells like those shown on the previous page.

MAIN IDEA TEKS 5A

The cell cycle has four main stages.

Just as all species have life cycles, from tiny chihuahuas to massive beluga whales, cells also have a life cycle. The **cell cycle** is the regular pattern of growth, DNA duplication, and cell division that occurs in eukaryotic cells. **FIGURE 1.1** shows its four main stages: gap 1, synthesis, gap 2, and mitosis. Gap 1, synthesis, and gap 2 together make up what is called interphase.

The stages of the cell cycle get their names from early studies of cell division. Scientists' observations were limited by the micro-


scopes of the time. When a cell was not actively dividing, they could not see activity in it. Thus, they

originally divided the cell cycle into two parts: interphase, when the cell appeared to be at rest,

and mitosis, when the cell was dividing.

Improved techniques and tools later allowed scientists to detect the copying of DNA (DNA synthesis), and they changed their description of the cell cycle to include the synthesis stage. Since they still could not see anything happening during the other parts of interphase, scientists named the periods between mitosis and synthesis "gap 1" and "gap 2." Eventually scientists learned that, during interphase, cells carry out their normal functions and undergo critical growth and preparation for cell division.

FIGURE 1.1 Cells grow and copy their DNA during interphase. They also carry out cell-specific functions in G_1 and G_2 . During M stage, both the nucleus (in mitosis) and cytoplasm (in cytokinesis) are divided.



INTERPHASE

Gap 1 (G₁)

The first stage of the cell cycle is gap 1 (G_1) . During G_1 , a cell carries out its normal functions. If it is a skeletal muscle cell, it contracts to move joints. If it is an adrenal cell, it secretes hormones such as adrenaline. If it is an intestinal cell, it absorbs nutrients. During G₁, cells also increase in size, and organelles increase in number. A cell spends most of its time in the G₁ stage, although the length of this stage varies by cell type.

During G₁, the cell must pass a critical checkpoint before it can proceed to the synthesis stage. Just as it would be dangerous for you to run a marathon if you had not slept or eaten for several days, it would also be dangerous for your cells to continue dividing if certain conditions were not met. For instance, most animal cells need enough nutrition, adequate size, and relatively undamaged DNA to divide successfully. They also need specific signals from other cells, telling them whether more cell division is needed.

Synthesis (S)


The second stage of the cell cycle is the synthesis (S) stage. *Synthesis* means "the combining of parts to make a whole." During the S stage, the cell makes a copy of its nuclear DNA. In eukaryotes, DNA is located in the nucleus. During interphase, it is loosely organized and appears grainy in photographs. By the end of the S stage, the cell nucleus contains two complete sets of DNA.

Gap 2 (G₃)

Gap 2 (G_2) is the third stage of the cell cycle. During G_2 , cells continue to carry out their normal functions, and additional growth occurs. Like G₁, this stage includes a critical checkpoint. Everything must be in order—adequate cell size, undamaged DNA—before the cell goes through mitosis and division.

Mitosis (M)


Mitosis (M), the fourth stage of the cell cycle, includes two processes: mitosis and cytokinesis. Mitosis (my-TOH-sihs) is the division of the cell nucleus and its contents. During mitosis, the nuclear membrane dissolves, the duplicated DNA condenses around proteins and separates, and two new nuclei form. Lastly, cytokinesis (sy-toh-kuh-NEE-sihs) is the process that divides the cell cytoplasm. The result is two daughter



cells that are genetically identical to the original cell.

The stages of the cell cycle and the proteins that control it are similar in all eukaryotes. For example, scientists have demonstrated that some of the molecules that regulate checkpoints in the yeast cell cycle can work in human cells, too. Such similarities suggest that eukaryotes share a common ancestry.

Predict What might happen if the G2 checkpoint stopped working in cells?



- CONNECT TO

DNA REPLICATION

As you will learn in the chapter From DNA to Proteins, DNA synthesis is also called DNA replication. During this process, the DNA molecule unzips and each strand is used as a pattern for a new DNA strand.



FIGURE 1.2 CELL LIFE SPAN **CELL TYPE APPROXIMATE LIFE SPAN** Skin cell 2 weeks Red blood cell 4 months Liver cell 300-500 days Intestine—internal 4-5 days lining Intestine—muscle and 16 years other tissues

Source: Spaulding et al., Cell 122:1.

CONNECT TO

LYMPHOCYTES

As you will learn in the chapter Immune System and Disease, lymphocytes are a part of your immune system. There are two major types of lymphocytes, B and T cells. Both types recognize specific antigens.

MAIN IDEA TEKS 5A

Cells divide at different rates.

Rates of cell division vary widely, as shown in **FIGURE 1.2**. The prokaryotic cell cycle is similar but not identical to that of eukaryotic cells. Recall that prokaryotes do not have the membrane-bound organelles and cytoskeleton found in eukaryotes. Thus, prokaryotic cells typically divide much faster than do eukaryotic cells.

The rate at which your cells divide is linked to your body's need for those cells. In human cells, the S, G₂, and M stages together usually take about 12 hours. The length of the G₁ stage differs most from cell type to cell type. The rate of cell division is greater in embryos and children than it is in adults. Children have a shorter cell cycle, and many of their organs are still developing. But the rate of cell division also varies within different tissues of the adult body. The internal lining of your digestive tract receives a lot of wear and tear. As a result, cells that line your stomach and intestine are replaced every few days. In contrast, cells that make up the rest of your intestine (mainly smooth muscle) and many of your internal organs, such as lungs,

kidneys, and liver, divide only occasionally, in response to injury or cell death.

Cells that divide only rarely are thought to enter a stage that some scientists call G_0 . In G_0 , cells are unlikely to divide, although they continue to carry out their normal functions. Some cells, such as neurons, appear to stay permanently in the G₀ stage. However, some data suggest that neurons actually can divide, and this question continues to be actively researched. Other cells, such as lymphocytes, a type of white blood cell, may remain in G_0 for years until they recognize an invader. Once the invader binds to a lymphocyte receptor, the lymphocyte goes through rapid cell divisions to help fight infection.

Infer Do you think a skin cell would have a long or short G₁ stage? Explain why.

MAIN IDEA

Cell size is limited.

Cells have upper and lower size limits. If cells were too small, they could not contain all of the necessary organelles and molecules. For instance, a cell with too few mitochondria would not have enough energy to live. However, cells cannot grow beyond a certain size, even if surrounded by plenty of nutrients. The upper limit on cell size is due to the ratio of cell surface area to volume. Recall that oxygen, nutrients, and wastes move across the cell membrane, or the surface of the cell. These materials must be transported in adequate amounts and with adequate speed to keep the inside of the cell functioning. But as a cell increases in size, its volume increases faster than its surface area, as shown in **FIGURE 1.3**. Therefore, a further increase in size could result in a surface area too small for the adequate exchange of materials.

FIGURE 1.3 Ratio of Surface Area to Volume in Cells

As a cell grows, its volume increases more rapidly than its surface area. When the surface area-to-volume ratio is too small, the cell cannot move materials into and out of the cell at a sufficient rate or in sufficient quantities.

Relative size	1-[2 —	3-	
Surface area (length \times width \times number of sides)	6	24	54	
	1	8	27	
Ratio of surface area to volume	6 1 = 6:1	24 /8 = 3:1	$\frac{54}{27} = 2:1$	

Compare Which cell has the largest surface area? Which cell has the largest surface area-to-volume ratio?

Some cells, however, must be large. A neuron running down a giraffe's neck to its legs may be several meters long, for instance. But it is not shaped like a cube or a sphere. Instead, it is extremely long and thin. This structure gives the neuron a large surface area with a relatively small increase in volume.

To maintain a suitable cell size, growth and division must be coordinated. If a cell more than doubled its size before dividing, the daughter cells would be larger than the original cell. If this happened with each generation, cells would quickly become too large to live. Similarly, if a cell did not double its size before dividing, the daughter cells would be smaller than the original cell. If this happened with each generation, cells would become too small to live.

Connect Which has the larger ratio of surface area to volume, a tennis ball or a soccer ball? Explain your reasoning.

Formative Assessment

REVIEWING D MAIN IDEAS

- 1. During which stage of the cell cycle is the DNA copied? TEKS 5A
- 2. Which stages of the cell cycle generally require about the same amount of time in all human cells? TEKS 5A
- 3. What limits the maximum size of a cell?

CRITICAL THINKING

- **4.** Infer Suppose you were to draw a diagram representing the cell cycle of a neuron. Explain where and how you would represent G₀. TEKS 5A
- **5. Predict** Suppose you treat cells with chemicals that block cytokinesis. Describe what you think the cells would look like.



CONNECT TO

SCIENTIFIC PROCESS

6. Predict how the rate of cell division would differ between single-celled algae living in a sunny, nutrient-rich pond versus algae living in a shady, nutrient-poor pond. How could you test your prediction?



5.2 Mitosis and Cytokinesis

TEKS 2G, 5A

VOCABULARY

chromosome histone chromatin chromatid centromere telomere prophase metaphase anaphase telophase

TEKS **2G,** analyze, evaluate, make inferences, and predict trends from data and 5A describe the stages of the cell cycle, including deoxyribonucleic acid (DNA) replication and mitosis, and the importance of the cell cycle to the growth of organisms

KEY CONCEPT Cells divide during mitosis and cytokinesis.

MAIN IDEAS

- Chromosomes condense at the start of mitosis.
- Mitosis and cytokinesis produce two genetically identical daughter cells.

- Connect to Your World

When you were a child, perhaps you attended a birthday party where goody bags were handed out. Whoever stuffed the bags had to make sure that each bag had exactly the same number of erasers, candies, and stickers. Otherwise, some illmannered child (not you, of course) might have raised a fuss if an item was missing. In a similar way, your cells must receive a full set of DNA—no more, no less—to work properly. Dividing DNA is a complicated task because the DNA is so long and stringy. Mitosis is an amazing process that efficiently sorts two sets of DNA and divides them between two nuclei.

MAIN IDEA TEKS 5A

Chromosomes condense at the start of mitosis.

DNA is a double-stranded molecule made of four different subunits called nucleotides. A **chromosome** is one long continuous thread of DNA that consists of numerous genes along with regulatory information. Your body cells have 46 chromosomes each. If stretched out straight and laid end to end, the DNA in just one of your cells would be about 3 meters (10 feet) long. How does it fit inside the nucleus of a microscopic cell?

DNA wraps around proteins that help organize and condense it. During interphase, or when a cell is not dividing, DNA is loosely organized—it looks a bit like spaghetti. During mitosis, however, your chromosomes are tightly condensed, as shown in **FIGURE 2.1**. These changes in DNA's organization allow a cell to carry out its necessary functions. During all of interphase, proteins must access specific genes for a cell to make specific proteins or to copy the entire DNA sequence. During mitosis, the duplicated chromosomes must condense to be divided between two nuclei.




FIGURE 2.1 This duplicated chromosome is tightly condensed. (colored SEM; magnification unknown)

If chromosomes remained stringy during mitosis, they could become entangled. Perhaps a cell would get two copies of one chromosome and no copies of a different one. **FIGURE 2.2** shows the process that converts a chromosome from a linear strand of DNA to its highly condensed form. The key to this process is the association between DNA and proteins.

CONNECT TO

BIOCHEMISTRY

As you will learn in the chapter From DNA to Proteins, a nucleotide is made of three parts: a sugar, a phosphate group, and a nitrogen-containing molecule called a base. When the sugars and phosphate groups bond, they form the backbones of the long chains called nucleic acids.



chromatid FIGURE 2.2 Chromosome Structure telomere DNA condenses tightly during the early stages of mitosis. centromere **DNA double helix** Condensed, duplicated chromosome **DNA** and histones Chromatin **Supercoiled DNA** Each continuous. Interactions between The chromatin The condensed, duplicated chromo-DNA wraps at regular double-stranded intervals around proparts of the histones coils more and more somes can be aligned and separated DNA molecule makes teins called histones, further compact the tightly around orgaduring mitosis. forming chromatin. one chromosome. DNA. nizing proteins. Infer Overall, DNA has a negative charge. Look at the histone proteins in the

At almost all times during the cell cycle, each of your chromosomes is associated with a group of proteins called **histones**. DNA wraps around histones at regular intervals, similar to beads on a string. The complex of protein and DNA that makes up the chromosome is called **chromatin**. Parts of the histones interact with each other, further compacting the DNA. At the "spaghetti" stage, the combination of DNA and proteins is loose. The word "loose" describes how much the DNA strand folds back on itself; it does not mean the DNA is loosely wrapped around the histones.

figure. What type of overall charge do you think they have? Explain.

As a cell progresses into mitosis, chromatin further condenses. It continues to coil more and more tightly around organizing proteins, finally forming small, thick rods. Recall that each chromosome has already been copied during the previous S stage. Thus, the chromosome looks similar to an "X" in which the left and right halves are two identical DNA double helixes. One half of a duplicated chromosome is called a **chromatid** (KROH-muh-tihd). Together, the two identical chromatids are called sister chromatids. Sister chromatids are held together at the centromere (SEHN-truh-MEER), a region of the condensed chromosome that looks pinched.

In addition, the ends of DNA molecules form structures called **telomeres** (TEHL-uh-MEERS), which are made of repeating nucleotides that do not form genes. They prevent the ends of chromosomes from accidentally attaching to each other, and they help prevent the loss of genes. A short section of nucleotides is lost from a new DNA molecule each time it is copied. It is important that these nucleotides are lost from telomeres, not from the genes themselves.

READING TOOLBOX TAKING NOTES Use a main idea web to help you study the makeup and organization of chromosomes. chromosomes


Apply What is the relationship between a molecule of DNA and a chromosome?

MAIN IDEA TEKS 5A

Mitosis and cytokinesis produce two genetically identical daughter cells.

The combined processes of mitosis and cytokinesis produce two genetically identical daughter cells. Follow along in FIGURE 2.4 as you read about the process in more detail below.

FIGURE 2.3 The nucleus and chromosomes go through dramatic changes in a dividing cell.



CONNECT TO

CELLS

As you will learn in the chapter Meiosis and Mendel, your body has two major cell types. Germ cells develop into eggs or sperm. Somatic cells make up the rest of vour body.



Interphase

Interphase plays an important role in preparing the cell to divide. It provides critical time for the duplication of organelles and for DNA replication. By the end of interphase, an individual cell has two full sets of DNA, or chromosomes, and is large enough to divide.





Mitosis **MITOSIS**

Mitosis divides a cell's nucleus into two genetically identical nuclei, each with its own single, full set of DNA. This process occurs in all of your body cells—except those that form eggs or sperm—and prepares them for cytokinesis. Although mitosis and cytokinesis are continuous processes, scientists have divided them into phases to make them easier to understand and discuss. The four main phases of mitosis are prophase, metaphase, anaphase, and telophase. Cytokinesis begins during late anaphase or telophase.

- During **prophase**, chromatin condenses into tightly coiled chromosomes. Each consists of two identical sister chromatids. The nuclear envelope breaks down, the nucleolus disappears, and the centrosomes and centrioles begin to migrate to opposite sides of the cell. Organized microtubules called spindle fibers grow from the centrioles and radiate toward the center of the cell.
- In metaphase, the spindle fibers attach to a protein structure on the centromere of each chromosome and align the chromosomes along the cell equator, around the middle of the cell.
- During **anaphase**, sister chromatids separate from each other. The spindle fibers begin to shorten, pulling the sister chromatids away from each other and toward opposite sides of the cell.
- 4 In **telophase**, a complete set of identical chromosomes is positioned at each pole of the cell. The nuclear membranes start to form, the chromosomes begin to uncoil, and the spindle fibers fall apart.

Cytokinesis

Cytokinesis divides the cytoplasm into two cells and completes a full stage of the cell cycle. Cytokinesis differs in animal and plant cells. In animal cells, the membrane forms a furrow, or trench, that is pulled inward by tiny filaments, like a drawstring. Gradually, the membrane pinches closed, forming a separate cell around each nucleus.

CYTOKINESIS



FIGURE 2.4 The Cell Cycle in Detail


Following interphase, mitosis divides duplicated chromosomes between two nuclei. Cytokinesis divides the cytoplasm. In this diagram, the mitosis stage is greatly **expanded to highlight its four major phases.** (micrographs; magnification about $100 \times$)

INTERPHASE


The cell copies its DNA and grows in preparation for division. The DNA is loosely organized during interphase.

MITOSIS

Mitosis divides a cell's nucleus into two nuclei, each with an identical set of DNA.



Prophase DNA and proteins condense into tightly coiled chromosomes. The nuclear envelope breaks down, centrioles begin to move to opposite poles, and spindle fibers form.



Metaphase Spindle fibers attach to each chromosome. They align the chromosomes along the cell equator.





Telophase Nuclear membranes start to form, chromosomes begin to uncoil, and the spindle fibers fall apart.

Anaphase Chromatids separate to opposite sides of the cell. Cytokinesis usually begins in late anaphase or telophase.



How many chromosomes does the cell have at the start of mitosis? How many does it have after cytokinesis? TEKS 5A

CYTOKINESIS

Cytokinesis divides cytoplasm between two daughter cells, each with

a genetically identical

interphase and begin

the cycle again.

nucleus. The cells enter

DATA ANALYSIS

TEKS 2G

CONSTRUCTING DATA TABLES

Scientists use data tables to record their data. Data tables are organized by the independent and dependent variables. Usually the independent variable is listed in the left column, and the other columns list the dependent variables. Each separate observation is listed in its own row. When measurements are taken using units, they are listed in the column headings in parentheses. All tables should have numbers and titles.

	TABLE 1. EFFECT OF HO	_			
	Concentration of Hormone Solution (%)	Size of Cell Clump After 24 Hours (mm)			
independent	0	3	dependent		
variable	25	4	variable		
	50	8			
	75	9			
	100	9			
observations					

Table 1 shows data from a hypothetical experiment in which growth hormones were added to clumps of cells in a laboratory, and the growth of the cell clumps was measured.

- 1. Display Data Suppose a scientist decided to measure the effect of temperature on cell division in chlorella, a type of green algae. Set up a table that could be used to record the results for the number of daily doublings of the cells: 20°C, 3 doublings; 30°C, 7 doublings; 40°C, 12 doublings; 50°C, 0 doublings.
- 2. Apply Label the independent and dependent variables on your table.



During cytokinesis in plant cells, the membrane cannot pinch inward because of the cell wall. Instead, a cell plate forms between the two nuclei. It is made by the Golgi apparatus, which supplies the new plasma membrane. A new wall then grows as cellulose and other materials are laid down. Typically, cytoplasm is divided evenly between daughter cells in both plant and animal cells.

The formation of new cells is critical in both multicellular and single-celled organisms. Single-celled organisms use cell division to reproduce, whereas multicellular organisms use it for growth, development, and repair.

Contrast How does cytokinesis differ in animal and plant cells?

Formative Assessment

REVIEWING A MAIN IDEAS

- 1. Draw what a **chromosome** looks like during **metaphase**. Identify the chromatids and the centromere.
- 2. Briefly explain why the daughter cells resulting from mitosis are genetically identical to each other and to the original cell. TEKS 5A

CRITICAL THINKING

- 3. Contrast How do prophase and telophase differ? TEKS 5A
- **4.** Apply Using a light microscope, you observe a cell that has no nucleus. What features would you look for to determine whether it is a eukaryotic cell undergoing mitosis or a prokaryotic cell? TEKS 5A

SELF-CHECK Online HMDScience.com

CONNECT TO

PROTEIN SYNTHESIS

5. For a cell to make proteins, enzymes must access its genes. When **histones** are modified with acetyl groups (-COCH₃), their positive charge is neutralized, so they wrap DNA less tightly. How might this affect the rate of protein synthesis? TEKS 4B





research.



Regulation of the Cell Cycle



5A, 5B, 5C, 5D, 9C

VOCABULARY

growth factor apoptosis cancer benign malignant metastasize carcinogen


TEKS 5A describe the stages of the cell cycle, including deoxyribonucleic acid (DNA) replication and mitosis, and the importance of the cell cycle to the growth of organisms; **5B** examine specialized cells, including roots, stems, and leaves of plants; and animal cells such as blood, muscle, and epithelium; **5C** describe the roles of DNA, ribonucleic acid (RNA), and environmental factors in cell differentiation; **5D** recognize that disruptions of the cell cycle lead to diseases such as cancer; 9C identify and investigate the role of enzymes

FIGURE 3.1 Normal animal cells (top) respond to external factors and stop dividing when they touch each other. Cancer cells (bottom) fail to respond and form clumps.

Normal cell growth



Cancerous cell growth



KEY CONCEPT Cell cycle regulation is necessary for healthy growth.

MAIN IDEAS

- Internal and external factors regulate cell division.
- O Cell division is uncontrolled in cancer.

-: Connect to Your World

Have you ever watched a movie in which people play with the elements of nature? They might bring back dinosaurs or make a newfangled robot. And have you noticed that these movies are always scary? That's because things go out of control. The robots take over, or the dinosaurs start eating humans. If cell growth goes out of control in your body, the result can be even scarier. Cancer is uncontrolled cell growth and results from many factors that affect the cell cycle. So how does your body regulate all the millions of cell divisions happening in your body?

MAIN IDEA TEKS 5A, 5B, 5C, 9C

Internal and external factors regulate cell division.

Both external and internal factors regulate the cell cycle in eukaryotic cells. External factors come from outside the cell. They include messages from nearby cells and from distant parts of the organism's body.

Internal factors come from inside the cell and include several types of molecules found in the cytoplasm. Both types of factors work together to help your body control the process of cell division.

External Factors

External factors that help regulate the cell cycle include physical and chemical signals. One example of a physical signal is cell-to-cell contact. Most mammal cells grown in the laboratory form a single layer on the bottom of a culture dish, as shown in **FIGURE 3.1**. Once a cell touches other cells, it stops dividing. The exact reason for this phenomenon is unknown. One hypothesis is that receptors on neighboring cells bind to each other and cause the cells' cyto-skeletons to form structures that may block the signals that trigger growth.

Many cells also release chemical signals that tell other cells to grow. For example, **growth factors** are a broad group of proteins that stimulate cell division. Growth factors bind to receptors that activate specific genes to trigger cell growth. In general, cells grow and divide in response to a combination of different growth factors, not just one.

Some growth factors affect many types of cells. For example, platelets are sticky fragments of bone marrow cells. They form clots that help stop bleeding. Platelets store a type of growth factor that helps your body repair wounds by triggering the growth of many cell types. Other growth factors have more specific targets. For instance, erythropoietin (ih-RIHTH-roh-poy-EE-tihn) stimulates the production only of cells that will become red blood cells. Red blood cells carry oxygen. If you moved from the coast to the mountains, your blood oxygen levels would be lower because the air pressure is lower at higher altitudes. The decrease in blood oxygen levels would cause your body to produce more erythropoietin. That factor would increase the number of red blood cells and raise your blood oxygen levels.

Various hormones may also stimulate the growth of certain cell types. In particular, growth hormone results in bone growth and also affects your protein and fat metabolism.

Internal Factors

When external factors bind to their receptors, they can trigger internal factors that affect the cell cycle. Two of the most important and well-studied internal factors involved in the eukaryotic cell cycle are kinases and cyclins. A kinase is an enzyme that, when activated, transfers a phosphate group from one molecule to a specific target molecule. This action typically increases the


energy of the target molecule or changes its shape. Your cells have many types of kinases, and they are almost always present in the cell. Those kinases that help control the cell cycle are activated by cyclins. Cyclins are a group of proteins that are rapidly made and destroyed at certain points in the cell cycle. These two factors help a cell advance to different stages of the cell cycle when cells bind to each other.

Apoptosis

Just as some cells need to grow and divide, other cells need to die. Apoptosis (AP-uhp-TOH-sihs) is programmed cell death. It occurs when internal or external signals activate genes that help produce self-destructive enzymes. Many questions remain about this process. What is known is that the nucleus of an apoptotic cell tends to shrink and break apart, and the cell is recognized by specialized cells in the immune system. These cells very tidily gobble up the apoptotic cell and recycle its chemical parts for use in building other molecules. **FIGURE 3.2** shows a classic example of apoptosis. In the early stages of development, human embryos have webbing between their fingers and toes, or digits. Before a baby is born, those cells typically go through apoptosis. Most babies are born with little unwebbed fingers and toes they love to put in their mouths.

Predict Suppose a child was born with growth hormone receptors that did not work properly. How do you think this would affect the child's development? TEKS 5A, 5C

FIGURE 3.2 Human embryos have webbed digits early in their development. The cells between the digits undergo apoptosis during later stages of development. As a result, the baby is born with unwebbed fingers and toes.







MAIN IDEA TEKS 5A, 5D

Cell division is uncontrolled in cancer.


FIGURE 3.3 Cancer cells form tumors that may metastasize to other parts of the body.



1. A healthy cell may become a cancer cell if certain genes are damaged.



2. Cancer cells divide more often than do healthy cells and may form disorganized clumps called tumors.



3. Sometimes, cancer cells break away from the tumor. They can be carried in the bloodstream to other parts of the body, where they form new tumors.

Cancer is the common name for a class of diseases characterized by uncontrolled cell division. It arises when regulation of the cell cycle is disrupted. Unlike healthy cells, cancer cells grown in a culture dish continue to divide, even when surrounded by neighboring cells. Cancer cells can also continue to divide in the absence of many of the growth factors required for division in healthy cells. As a result, they divide much more often than do healthy cells.

Cancer cells form disorganized clumps called tumors. In a benign tumor, the cancer cells typically remain clustered together. This means the tumor may be relatively harmless and can probably be cured by removing it. However, if a tumor is malignant, some of the cancer cells can break away, or metastasize (mih-TAS-tuh-syz), from the tumor. These breakaway cells can be carried in the bloodstream or lymphatic system to other parts of the body, as shown in FIGURE 3.3, where they can form more tumors, called metastases. Once a tumor metastasizes, it is much more difficult to entirely rid the body of tumors.

But why are tumors harmful? Cancer cells do not perform the specialized functions needed by the body. In the lung, for example, cancer cells do not exchange oxygen and carbon dioxide. In the brain, they do not transmit the carefully ordered electrical messages needed to interpret information. Therefore, the body has large clumps of rapidly dividing cells that require lots of food and a hearty blood supply but that contribute nothing to the body's function. In addition, a growing tumor can exert great pressure on surrounding organs. For instance, a tumor growing inside the skull will cramp the brain for space, and some regions will be unable to function properly. If cancer cells continue to grow unchecked, they will eventually kill the organism.

Cancer cells come from normal cells that have suffered damage to the genes that help make proteins involved in cell-cycle regulation. Most cancer cells carry mutations, or errors, in two types of genes. One type, called oncogenes, accelerate the cell cycle. The second type act as cell-cycle brakes. Mutations in these genes can be inherited. For instance, some breast

cancers appear to be caused by inherited errors in specific genes. Other mutations can be caused by exposure to radiation or chemicals. For example, some skin cancers are due to DNA damage caused by ultraviolet radiation from sunlight. Substances known to produce or promote the development of cancer are called carcinogens (kahr-SIHN-uh-juhnz). These include tobacco smoke and certain air pollutants, which are both associated with lung cancer. Some mutated forms of oncogenes are even carried by viruses; one such virus can cause cervical cancer.




FIGURE 3.4 This cancerous mole is an example of a skin cancer, which may metastasize quickly.

QUICK LAB

OBSERVING TEKS 5B

Cancer

In this lab, you will compare normal cells with cancerous cells and observe the differences between them.

PROBLEM How do normal and cancerous cells compare?

PROCEDURE

- 1. Examine the slides of normal cells under the microscope. Draw and describe your observations.
- 2. Repeat step 1 with slides of cancer cells.

ANALYZE AND CONCLUDE

- 1. Compare How does the structure of the normal cells compare with the structure of the cancerous cells for each of the slides you viewed?
- 2. Infer Cancer cells not only appear different from normal cells but they also divide more rapidly. Why do you think chemotherapy, a common treatment for cancer, results in the loss of hair?

MATERIALS

- microscope
- slides of normal cells
- slides of cancerous cells

Standard cancer treatment often involves both radiation and chemotherapy. Radiation therapy is the use of radiation to kill cancer cells and shrink tumors. It works by damaging a cell's DNA so much that the cell cannot divide. Radiation is usually localized—that is, its use is targeted to a specific region—because it can also hurt healthy cells. Chemotherapy uses certain drugs, often in combination, to kill actively dividing cells. Like radiation, it kills both cancerous and healthy cells. However, chemotherapy is systemic drugs travel throughout the entire body.

Medical researchers use laboratory-grown cancer cells in their search for cancer treatments. Much of what is known about the cell cycle has come from studies that use cancer cells. The most famous cancer cells used for research are called HeLa cells. HeLa cells were originally obtained in 1951 from a cervical tumor removed from a woman named Henrietta Lacks. This cell line continues to be grown and studied in laboratories all over the world.

Analyze HeLa cells are also used to study cell signaling processes. What might be a disadvantage of using cancer cells to study processes occurring in healthy cells? TEKS 5D

Web*Ques*t HMDScience.com Skin Cancer

Formative Assessment

REVIEWING MAIN IDEAS

- 1. Describe what a growth factor is and how it influences the cell cycle. TEKS 5A
- 2. Explain how cancer cells differ from healthy cells. **TEKS 5A, 5D**

CRITICAL THINKING

- 3. Contrast How do benign and malignant tumors differ?
- **4. Hypothesize** Suppose chromosomes in a skin cell are damaged by ultraviolet radiation. If the damaged genes do not affect cell cycle regulation, do you think the cell will become cancerous? Explain. TEKS 5A, 5D



CONNECT TO

CELL ORGANELLES

5. Some anticancer drugs prevent microtubules from forming spindle fibers. Why do you think these drugs might be effective treatments for cancer? TEKS 5A, 5D

5.4 Asexual Reproduction

VOCABULARY

asexual reproduction binary fission

KEY CONCEPT Many organisms reproduce by cell division.

MAIN IDEAS

- Dinary fission is similar in function to mitosis.
- Some eukaryotes reproduce through mitosis.

- Connect to Your World

In this flashy world of ours, you may think that the humble bacterium would have little chance of finding a mate. No dazzling smile, no fancy hair products, no shiny car, and—if we are brutally honest—not even a brain. With all of these limitations, it may seem that our bacteria friends would be destined to die out. And yet, bacteria are found in abundance and live just about everywhere on Earth. How can there be so many bacteria?

MAIN IDEA

Binary fission is similar in function to mitosis.

Reproduction is a process that makes new organisms from one or more parent organisms. It happens in two ways—sexually and asexually. Sexual reproduction involves the joining of two specialized cells called gametes (eggs and sperm cells), one from each of two parents. The offspring that result are genetically unique; they have a mixture of genes from both parents. In contrast, **asexual reproduction** is the production of offspring from a single parent and does not involve the joining of gametes. The offspring that result are, for the most part, genetically identical to each other and to the single parent.

nuclei. They also do not have spindle fibers. And although they have DNA, prokaryotes have much less DNA than do most eukaryotes. The DNA of most

bacteria is in the form of a single circular chromosome.

Binary Fission and Mitosis

Most prokaryotes reproduce through binary fission (BY-nuh-ree FIHSHuhn), the asexual reproduction of a single-celled organism by which the cell divides into two cells of the same size. Binary fission and mitosis have similar results. That is, both processes form two daughter cells that are genetically identical to the parent cell. However, the actual processes are different in several important ways.

As you already learned, prokaryotes such as bacteria do not have

VISUAL VOCAB Binary fission is the asexual reproduction of a single-celled organism by division into two roughly equal parts. parent cell **DNA** duplicates cell begins to divide daughter cells

CONNECT TO

CELL STRUCTURE

Recall from the chapter Cell Structure and Function that many scientists hypothesize that mitochondria and chloroplasts were originally free-living prokaryotes. One piece of evidence that supports this hypothesis is the fact that these two organelles replicate much as bacteria do, through fission.




FIGURE 4.1 This micrograph shows three individual bacteria, each at a different stage of binary fission. First, a cell elongates (1), and the DNA is replicated. Next, the cell membrane pinches inward (2). Finally, the membrane meets, and a new cell wall forms, separating the two cells (3). (colored TEM; magnification 60,000×)


Binary fission, shown in **FIGURE 4.1**, starts when the bacterial chromosome is copied. Both chromosomes are attached to the cell membrane. As the cell grows and gets longer, the chromosomes move away from each other. When the cell is about twice its original size, it undergoes cytokinesis. The membrane pinches inward, and a new cell wall forms between the two chromosomes, which completes the separation into two daughter cells.

Advantages and Disadvantages of Asexual Reproduction

Very often, whether something is helpful or harmful depends on the situation. In favorable environments that do not change much, asexual reproduction can be more efficient than sexual reproduction. Recall that asexual reproduction results in genetically identical offspring. If they are well suited to the environment, genetic variation could be more harmful than helpful. In other words, if it ain't broke, don't fix it.

However, asexual reproduction may be a disadvantage in changing conditions. Genetically identical offspring will respond to the environment in the same way. If population members lack traits that enable them to reproduce in a changed environment, the entire population could die off. In contrast, sexual reproduction increases genetic diversity, which raises the chance that some individuals will survive in changing conditions.

Keep in mind, however, that the act of asexual reproduction itself is not more efficient; rather, the associated costs of sexual reproduction are greater. For example, all asexually reproducing organisms can potentially reproduce. Suppose two organisms each have ten offspring. If one organism reproduces asexually, all ten offspring can have offspring of their own. If the other organism reproduces sexually, having five females and five males, only the five females can bear offspring. In addition, sexually reproducing organisms must attract a mate. This effort involves not only the time and energy needed to find a mate but also many structures, signals, and behaviors that have evolved to attract mates. Organisms that reproduce asexually do not have these costs.



of 1024 cells after only 10 rounds of cell division.

CONNECT TO

EVOLUTION

As you will learn in the chapter Viruses and Prokaryotes, the misuse of antibiotics has resulted in multidrug-resistant bacteria. The bacteria not killed by antibiotics can reproduce quickly, passing the genes for antibiotic resistance on to their offspring.

Summarize How is asexual reproduction an advantage in some conditions?

Some eukaryotes reproduce through mitosis.

FIGURE 4.3 Yeast and hydras can reproduce by budding. (hydra: LM, magnification 12×; yeast: colored SEM, magnification 3,200×)



Some eukaryotes also reproduce asexually, through mitosis. Have you ever grown a new plant from a stem cutting? Or seen a new sea star growing from the arm of another one? These new organisms are the result of mitotic reproduction and are therefore genetically the same as the parent organism. Mitotic reproduction is especially common in simpler plants and animals. It occurs in both multicellular and unicellular eukaryotes. It can take several forms, including budding, fragmentation, and vegetative reproduction.

In budding, a small projection grows on the surface of the parent organism, forming a separate new individual. The new organism may live independently or attached as part of a colony. For instance, hydras and some types of yeast reproduce by budding. Examples are shown in **FIGURE 4.3**.

In fragmentation, a parent organism splits into pieces, each of which can grow into a new organism. Flatworms and sea stars both reproduce by fragmentation. Many plants, including strawberries and potatoes, reproduce via vegetative reproduction. In general, vegetative reproduction involves the modification of a stem or underground structures of the parent organism. The offspring often stay connected to the original organism, through structures called runners, for example.

Many organisms can reproduce both asexually and sexually. The form of reproduction may depend on the current conditions. The sea anemone can reproduce in many ways. It can reproduce asexually by dividing in half, by breaking off small pieces from its base, or by budding. It can also reproduce sexually by making eggs and sperm. Some species of anemone have separate males and females. In other anemone species, the same organism can produce both eggs and sperm cells.

Synthesize How might the asexual reproduction of genetically identical plants be useful to humans? How could it prove harmful to our food supply?

5.4 Formative Assessment

REVIEWING MAIN IDEAS

- **1.** Explain how mitosis differs from **binary fission.**
- **2.** Briefly explain why cutting a flatworm into pieces would not kill it.

CRITICAL THINKING

- 3. Infer How does an organism benefit by being able to reproduce both sexually and asexually?
- **4. Apply** Yeasts are growing in two dishes. You treat one dish with a chemical that blocks DNA replication but forget to label it. How can you identify the treated dish?

SELF-CHECK Online HMDScience.com

CONNECT TO

ECOLOGY

5. Two populations live in the same habitat and compete for food. The first group is larger and multiplies through asexual reproduction; the second reproduces sexually. What could happen to cause the second group to outnumber the first?

(t)) @Rotandi Birke/Phototake Inc./Alamy Ltd; (b) @J. Forsdyke/Gene Cox/Photo Researchers, Inc.



Multicellular Life



5B, 5C, 10C

VOCABULARY

tissue organ organ system cell differentiation stem cell

TEKS 5B examine specialized cells, including roots, stems, and leaves of plants; and animal cells such as blood, muscle, and epithelium; **5C** describe the roles of DNA, ribonucleic acid (RNA), and environmental factors in cell differentiation; 10C analyze the levels of organization in biological systems and relate the levels to each other and to the whole system

KEY CONCEPT Cells work together to carry out complex functions.

MAIN IDEAS

- Multicellular organisms depend on interactions among different cell types.
- Specialized cells perform specific functions.
- Stem cells can develop into different cell types.

-: Connect to Your World

Each of us enters this world as a helpless infant. At first, your ability to eat solid foods or take your first steps elicits a great deal of praise. Over time, however, your development of normal skills gets far less attention. By the time you reach the age of 18, people want to know what you plan to do with your life. Will you build houses or design clothing or treat patients? What will your specialty be? Cells, too, undergo specialization to carry out the complex functions required by the body.

MAIN IDEA TEKS 10C

Multicellular organisms depend on interactions among different cell types.

Within multicellular organisms, cells communicate and work together in groups that form increasingly larger, more complex structures. This arrangement progresses from cells to tissues to organs to organ systems, as shown in **FIGURE 5.1.** Tissues are groups of cells that work together to perform a similar function. Groups of tissues that work together to perform a specific function or related functions are called **organs**. For instance, plants have photosynthetic tissues made of chlorophyll-containing cells. Conductive tissues transport sugars, water, and minerals to and from other parts of the plant. Protective tissues help prevent water loss. Together, these and other tissues form a leaf, the plant's food-producing organ.

Organs that carry out similar functions are further grouped into **organ systems.** In plants, the shoot system is above the ground. It includes stems that support the plant, leaves that capture radiant energy, and flowers that aid reproduction. Beneath the ground, the root system has different types of roots and root hairs that anchor the plant and absorb water and minerals.

As organ systems work together, they help an organism maintain homeostasis. For example, plants need to maintain a certain level of water within their cells, otherwise they will wilt and die. They absorb water through their roots and expel it as water vapor through openings in their leaves called stomata. Stomata are controlled by special cells called guard cells, which close the stomata when a plant's water intake cannot keep up with its water loss.

Apply Suppose your family goes out of town and forgets to ask your neighbor to water the plants. Do you think the plants' stomata will be open or closed? Explain.

CONNECT TO

HOMEOSTASIS

As you learned in the chapter Biology in the 21st Century, homeostasis is the maintenance of a stable internal environment. Both an organism's physiology and its behavior help it achieve homeostasis.

FIGURE 5.1 Levels of Organization

Cells work together in groups that form larger, specialized structures.

CELL

Vessel elements are tube-shaped cells. (colored SEM; magnification 200×)




TISSUE


Vessel elements, tracheids, and parenchyma cells form xylem. (colored SEM; magnification 240×)



ORGAN

Xylem and other tissues form roots that absorb water and nutrients.





Apply How is the shape of this plant's roots suited to their function? TEKS 5B, 10c

MAIN IDEA TEKS 5B, 5C

Specialized cells perform specific functions.

CONNECT TO

GAMETOGENESIS

As you will learn in the chapter **Meiosis and Mendel**, the egg is stocked with organelles and molecules that are necessary for an embryo to grow. Many of these molecules are not evenly distributed throughout the cell; they form gradients.


It is easy to see that a skin cell can divide to make a new skin cell, or that a single bacterium can generate another bacterium. But how does a complex organism like you develop? Your body began as a single fertilized egg. If the egg simply divided to make lots of identical cells, it would not form a baby. To form the intricate structures that make up your body and the bodies of countless organisms around you, cells must specialize.

Cell differentiation is the process by which a cell becomes specialized for a specific structure or function during multicellular development. While almost every cell in your body has a full set of DNA, each type of cell expresses only the specific genes it needs to carry out its function. That is, a cell differentiates among the genes and uses only certain ones. You can think of your DNA as a cookbook. When you want to make a specific dish, you select that recipe and carry out its instructions. If you need to make a dessert, you might choose brownies. If you need to make a main course, you might fix lentil stew. The dishes are very different, but they all come from the same cookbook.


A cell's location within the embryo helps determine how it will differentiate. In plant cells, the first division of a fertilized egg is unequal, or asymmetric, as shown in **FIGURE 5.2**. The apical cell forms most of the embryo, including the growth point for stems and leaves. The major role of the basal cell is to provide nutrients to the embryo; it also creates the growth point for the roots. Plant cells cannot easily migrate because of the cell wall, but they adapt to changing conditions and continue to develop throughout their lifetime.

As the plant grows, new cells continue to differentiate based on their location. For example, cells on the outer layer of a leaf may become epidermal cells that secrete a waxy substance that helps prevent water loss. Cells on the lower leaf surface may become guard cells that control the exchange of water, air, and carbon dioxide.

FIGURE 5.2 Cell Differentiation in Seed Plants



In animals, an egg undergoes many rapid divisions after it is fertilized. The resulting cells can migrate to a specific area, and the cells quickly begin to differentiate. The early animal embryo generally takes the shape of a hollow ball. As the embryo develops, part of the ball folds inward, forming an inner layer and creating an opening in the outer cell layer. A middle layer of cells then forms. As shown in FIGURE 5.3, in vertebrates, the outer cell layer differentiates to form the outer layer of skin and elements of the nervous system, such as the brain and spinal cord. The middle cell layer forms bones, muscles, kidneys, and the inner layer of skin. The inner cell layer forms internal organs, such as the pancreas, lungs, and digestive system lining.



CONNECT TO


VOCABULARY

Potent comes from a Latin word meaning "to be able."
The addition of prefixes defines the level of power or ability.
toti- = all
pluri- = more, several
multi- = many



Stem cells can develop into different cell types.

Stem cells are a unique type of body cell that can (1) divide and renew themselves for long periods of time, (2) remain undifferentiated in form, and (3) differentiate into a variety of specialized cell types. When a stem cell divides, it forms either two stem cells or one stem cell and one specialized cell.





Stem Cell Classification

Stem cells can be classified by their ability, or potential, to develop into the differentiated cell types of different tissues. In general, the more differentiated a stem cell already is, the fewer the types of cells it can form.

- Totipotent stem cells can grow into any other cell type. Only a fertilized egg and the cells produced by the first few divisions of an embryo are totipotent.
- Pluripotent stem cells can grow into any cell type except for totipotent stem cells.
- Multipotent stem cells can grow only into cells of a closely related cell family.

Stem cells are also classified by their origin, as either adult or embryonic. Adult stem cells have been studied for decades, but the ability to grow human embryonic stem cells was not developed until 1998. Since that time, embryonic stem cells have attracted great attention because of their potential to form almost any cell type.

Adult Stem Cells

Adult stem cells are partially undifferentiated cells located among the specialized cells of many organs and tissues. They are found all over the body, in the brain, liver, bone marrow, skeletal muscle, dental pulp, and even fat. These stem cells are also found in children and in umbilical cord blood, so the term *somatic stem cell* is more accurate although less frequently used.

A major advantage of adult stem cells is that they can be taken from a patient, grown in culture, and put back into the patient. Thus, the risk of transplant rejection by a patient's immune system is very low. This method also avoids many ethical issues associated with using embryonic stem cells.

Adult stem cells currently pose many disadvantages as well. They are few in number, difficult to isolate, and sometimes tricky to grow. They may also contain more DNA abnormalities than do embryonic stem cells. For years, much evidence suggested that adult stem cells were multipotent. This would mean that a stem cell from fat would produce only fat cells, never muscle cells.

Newer data suggest otherwise. Adult stem cells treated with the right combination of molecules may give rise to a completely different type of tissue. This process, called transdifferentiation, remains an active area of research.

Embryonic Stem Cells

Most embryonic stem cells come from donated embryos grown in a clinic. These embryos are the result of in vitro fertilization, a process by which eggs are fertilized outside a woman's body. The stem cells are taken from a cluster of undifferentiated cells in the three-to-five-day-old embryo. These cells,

called the inner cell mass, do not have the characteristics of any specific cell type. Because they are pluripotent, they can form any of the 200 cell types of the body. They can also be grown indefinitely in culture. These qualities offer hope that many diseases will be treatable or even curable. Stem cells have long been used to treat patients with leukemia and lymphoma, and people with diabetes might someday be cured if nonworking cells in the pancreas are replaced with healthy, growing cells. Even damaged organs might be strengthened by an injection of healthy cells.

Embryonic stem cells also have a downside. If these cells are used in treatment, a patient's body might reject them as foreign material. A different possibility is that the stem cells could grow unchecked in a patient's body and form a tumor. The use of embryonic stem cells also raises many ethical questions. FIGURE 5.4 shows the most common method of obtaining embryonic stem cells. This method currently

FIGURE 5.4 HARVESTING EMBRYONIC STEM CELLS inner cell mass fertilized egg muscle cells red blood cells neurons

First, an egg is fertilized by a sperm cell in a petri dish. The egg divides, forming an inner cell mass. These cells are then removed and grown with nutrients. Scientists try to control how the cells specialize by adding or removing certain molecules.

involves destruction of the embryo, which some people consider ethically unacceptable.

Compare and Contrast List treatment benefits and risks of both types of stem cells.

Formative Assessment

REVIEWING A MAIN IDEAS

- **1.** How does communication between cells help maintain homeostasis? TEKS 10C
- 2. Why is **cell differentiation** an important part of the development of a multicellular organism?
- **3.** What are the defining characteristics of stem cells?

CRITICAL THINKING

- **4. Compare** Describe how cells, tissues, organs, and organ systems are related. TEKS 100
- **5. Evaluate** What role does the location of a cell in a developing embryo play in cell differentiation? TEKS 5C

SELF-CHECK Online HMDScience.com

CONNECT TO

ETHICS IN BIOLOGY

6. Explain which factor you think is most important in deciding whether stem cell research should be legal and government-funded. TEKS 3D

Summary


BIG IDEA Cells have their own life cycle that includes reproduction, growth, and regulation, which allows organisms to carry out life functions and grow.

KEY CONCEPTS

5.1 The Cell Cycle

Cells have distinct phases of growth, reproduction, and normal functions. The cell cycle has four main stages: G_1 , S, G_2 , and M. The length of the cell cycle can vary, resulting in different rates of cell division. This variability is based on the body's need for

different cell types. Cells also divide because they need a sufficient surface area-to-volume ratio to move materials into and out of the cell.



5.2 Mitosis and Cytokinesis

Cells divide during mitosis and cytokinesis. Mitosis divides the nucleus into two genetically identical nuclei in a four-phase process: prophase, metaphase, anaphase, and telophase. In prophase, the duplicated chromosomes condense tightly. Cytokinesis actually divides the cell cytoplasm.

5.3 Regulation of the Cell Cycle


Cell cycle regulation is necessary for healthy growth. Cell growth and division are regulated by both external factors, such as hormones and growth factors, and internal factors, such as cyclins and kinases. When proper regulation of cell growth is disrupted, a cell may become cancerous. Cancer cells grow more rapidly than do normal cells and form clumps called tumors that may metastasize to other regions of the body.

5.4 Asexual Reproduction

Many organisms reproduce by cell division. Most prokaryotes reproduce through a process called binary fission, in which a cell divides into two approximately equal parts. Some eukaryotes reproduce through mitosis. The offspring that result from asexual reproduction are genetically identical to the parent organism, except when mutations occur. Whether being identical is an advantage or a disadvantage depends on the environment.

5.5 Multicellular Life


Cells work together to carry out complex functions. Within multicellular organisms, cells form tissues, tissues form organs, and organs form organ systems. The cells differentiate to perform specific functions. Much of this specialization is determined by a cell's location within the developing embryo. Stem cells are a special type of cell that continue to divide and renew themselves for long periods of time.




READING TOOLBOX

SYNTHESIZE YOUR NOTES

Concept Map Use a concept map like the one below to summarize what you know about mitosis.



Venn Diagram Draw a Venn diagram like the one below to summarize the similarities and differences between embryonic and adult stem cells.



Review Games • Concept Map • Section Self-Checks

CHAPTER VOCABULARY

- 5.1 cell cycle mitosis cytokinesis
- **5.2** chromosome histone chromatin chromatid centromere telomere


- prophase metaphase anaphase telophase
- growth factor apoptosis cancer benign malignant

- metastasize carcinogen
- 5.4 asexual reproduction binary fission
- 5.5 tissue organ organ system cell differentiation stem cell

Reviewing Vocabulary

Visualize Vocabulary

For each term below, draw a simple picture that represents the meaning of the word. Here is an example for mitosis.



- 1. prophase
- 2. metaphase
- 3. anaphase
- 4. telophase
- 5. cytokinesis
- 6. centromere
- 7. telomere

Reviewing MAIN IDEAS

- **11.** The cell cycle has four main stages— G_1 , S, G_2 , and M. What occurs in the cell during each stage? TEKS 5A
- 12. Compare the rates of cell division occurring in your neurons and your hair follicles. TEKS 5A
- 13. What is the relationship between a cell's surface area and its volume?
- **14.** You know that a chromosome is a very long, continuous strand of DNA. How do proteins help condense chromosomes?
- 15. Describe what happens in each main phase of mitosis prophase, metaphase, anaphase, and telophase. TEKS 5A
- 16. How does the process of cytokinesis differ from the process of mitosis? TEKS 5A
- 17. Increased levels of cyclin help trigger a cell to divide. Do you think a growth factor would increase or decrease cyclin levels? Explain. TEKS 9C
- 18. Describe how uncontrolled cell division is dangerous in organisms. TEKS 5D
- 19. List one similarity and one difference between binary fission and mitosis.
- 20. You pull a leaf from a plant and place it in a cup of water. After a week, roots start to grow from the leaf. What type of reproduction has occurred, and what role does mitosis play in it? TEKS 5A
- 21. Briefly describe how cell differentiation occurs in the developing animal embryo.
- 22. List three characteristics of all stem cells.

READING TOOLBOX WORD ORIGINS

- **8.** The prefix *pro-* means "earlier than" or "prior to." Explain how this meaning relates to the word *prophase*.
- **9.** The prefix *telo-* means "distant, far, or end." How does this meaning relate to the words telophase and telomere?
- **10.** The term *mitosis* comes from the Greek root *mitos*, which means "thread." How does this meaning relate to the process of mitosis?


Critical Thinking

- **23. Synthesize** How do regulatory proteins of the cell cycle help maintain homeostasis?
- **24. Describe** How is the location of a cell in an embryo related to differentiation of that cell? **TEKS 50**
- **25. Analyze** A scientist wants to use asexually reproducing vegetables to increase crop yields. He plans to distribute budding potatoes and teach farmers how to separate them into new plants. What are some potential benefits and risks that could result from this situation?
- **26. Analyze** The rates of DNA mutations in bacteria are known to increase when they are under stressed environmental conditions. Why do you think this is important for an organism that reproduces asexually?
- **27. Apply** Suppose an organism usually has 24 chromosomes in its nucleus. How many chromatids would it have just after the S phase of the cell cycle? **TEKS 5A**
- **28. Predict** If a mutation made histone proteins bind less tightly to DNA, how might the cell cycle be affected?

 TEKS 5A

Interpreting Visuals

Use the picture of onion root cells shown below to answer the next three questions.



magnification 120imes

- **29. Apply** In what stage of the cell cycle are most of these cells? Explain. **TEKS 5A, 5B**
- **30. Apply** How can you visually distinguish between newly formed cells and older cells? **TEKS 5A, 5B**
- **31. Synthesize** If these cells were immersed in salt water, how would the cells undergoing mitosis be affected? (Hint: Think about the process of osmosis.) **TEKS 5A**

Analyzing Data Construct a Data Table

The graph below shows the five-year survival rate, expressed as percentages, of patients diagnosed with cancer from 1985 through 1997. This data is for all types of invasive cancers and includes males and females of all races. Use the graph to answer the next two questions.

5-YEAR CANCER SURVIVAL RATE



Source: The National Cancer Institute

- **32. Analyze** Which points do not follow the best-fit line for the data?
- **33. Interpret** What is the trend in the data of cancer survival during the span of time given?

Making Connections

- 34. Write a Narrative Imagine that you are a single chromosome about to undergo replication and mitosis.

 Describe what will happen to you starting from the S phase through mitosis. Be creative. Use humor and a first-person point of view. Come up with sounds or perspectives that illustrate what is happening. Be sure to include all details of the process and related terms.
- 35. Design an Experiment Cancer cells, such as those shown on the chapter opener, are frequently grown in labs for research uses. Suppose you wanted to determine whether a certain substance was a carcinogen. Outline a plan for an experiment to describe what questions you would want to answer, what experiments you would perform, and what the different possible results would suggest. TEKS 5D



Biology End-of-Course Exam Practice

Record your answers on a separate piece of paper.

MULTIPLE CHOICE

TEKS 3A, 5D

- 1 Scientists researching anticancer drugs treat a cell culture with a compound. Following treatment, they notice that the culture stopped growing. Untreated cells from the same culture, however, have continued to grow. These results indicate that the compound blocks the normal cell cycle. What else could have caused these results?
 - A The compound had degraded.
 - B The compound prevented cells from mutating.
 - C The compound killed the treated cells.
 - D The compound had no effect.

THINK THROUGH THE QUESTION

The untreated cells serve as a control in this experiment. Therefore, differences between the treated and untreated cells should be the result of the drug. If the drug has no effect, the two groups of cells should be the same.


TEKS 5C

- 2 Which describes the role of DNA in cell differentiation?
 - A It regulates cell differentiation.
 - B It carries the code from which each cell type expresses specific genes.
 - C It removes genes that are unneeded from differentiated cells.
 - D It prevents a full set of DNA from being passed on to stem cells.

TEKS 5C

- 3 One environmental factor that plays a key role in cell differentiation in multicellular organisms is the location of the developing cell with respect to other cells. In human development, from which cell layer in the embryo will a nerve cell develop?
 - A within the nucleus
 - B outer layer
 - C middle layer
 - D inner layer


TEKS 4B, 5A



In the diagram above, cell A is undergoing mitosis. If cell A has 6 chromosomes, how many chromosomes will cells B and C have?

- A none
- B 3 each
- C 6 each
- D 12 each

TEKS 10C



The figure above represents some levels of organization in multicellular organisms. Which term fits in the box marked "X"?

- A cells
- B organelles
- C organs
- D organisms

TEKS 5C

- **6** Unlike stem cells, most body cells cannot form different types of cells. For example, skin cells can only make skin cells, and nerve cells only make nerve cells. Which statement **best** explains why skin cells will never become nerve cells?
 - A Each type of cell gets a different message from the central DNA, which is stored in DNA cells.
 - B Each type of cell has only the part of the DNA necessary for making that type of cell.
 - C Each cell type is determined by messages sent from the brain, which directs development.
 - D Both types of cells have the same DNA, but each cell uses only part of the DNA message.

at HMDSCIENCE.COM

INTERNET MAGAZINE

Go online for the latest biology news and updates on all BioZine articles.

Expanding the Textbook

News Feeds

- Science Daily
- **™** CNN
- **BBC**




Careers

Bio Bytes

Opinion Poll

Strange Biology



Stem Cell Research-Potential Solutions, **Practical Challenges**

A news program asks viewers to vote online: "Should stem cell research be banned? Yes or no?" Some people claim that stem cell therapy will revolutionize medicine. Others believe that some types of stem cell research violate ethical standards and are not justified by the potential benefits. Between these two positions exists a wide range of ideas about what is or is not acceptable. Would you know how to vote?

Using Stem Cells TEKS 3D

Stem cells are undifferentiated cells that can regenerate themselves and develop into specialized types of cells. Stem cell research offers the hope of understanding basic cell processes and treating or even curing many diseases. However, many technical challenges must be overcome before stem cell therapy is a realistic option, and ethical issues continue to surround stem cell research.

Potential Benefits TEKS 3D

Stem cell research offers many potential benefits.

 Studying adult stem cells may help scientists better understand how tissues develop and what goes wrong when those tissues become diseased.

- A better understanding of the properties of stem cells may give scientists more information about how cancer cells replace themselves and thus help scientists develop more-targeted cancer therapies.
- Stem cells could be used to grow human tissues to test the effects of drugs and chemicals.
- Stem cells may be used to replace healthy cells that are killed by radiation treatment for cancer.
- Stem cells may be used to regenerate tissues. For example, chemotherapy kills blood-producing cells in bone marrow. To replace these cells, stem cells could be used instead of the patient's own marrow, which may contain cancer cells.
- · Stem cells may be used to treat spinal cord injuries and neurodegenerative diseases, such as Parkinson's.

TECHNOLOGY **STEM**


Somatic Cell Nuclear Transfer

Somatic cell nuclear transfer (SCNT), also called therapeutic cloning, is a method for obtaining stem cells that has been used to clone animals. The process is still under development, however, and it has not yet been used to produce stem cells for humans. SCNT offers the hope of using a patient's own DNA to produce stem cells that can form many types of specialized cells. Many SCNT studies have been done in mice and pigs; the diagram to the right shows how the SCNT process might be applied in human cells.

- 1 An unfertilized egg is taken from a female's body, and the nucleus—containing the DNA—is removed. A cell is then taken from a patient's body. The nucleus is removed and inserted into the egg.
- The egg is given a mild electrical stimulation, which makes it divide. The DNA comes from the patient's nucleus, and the materials needed for division come from the egg.
- The stem cells could then be cultured and caused to differentiate into any tissue or organ needed by the patient.

Once a stem cell line is established, in theory it can continue to grow indefinitely. Researchers could use these cell lines without having to harvest more stem cells. The cell lines also could be frozen and shipped to other researchers around the world.

Read More >> at HMDScience.com



Technical Challenges

Adult stem cells have been used therapeutically for years in the form of bone marrow transplants. Nevertheless, many technical challenges must still be overcome before stem cells can be used to treat a wide range of disorders. Examples are highlighted below.

Supply Stem cells can be taken from a variety of sources, including an embryo, a patient in need of treatment, a patient's relative, or an established embryonic stem cell line. Embryonic stem cells are taken from embryos fertilized in an in vitro fertilization clinic, whereas established stem cell lines are cultures of embryonic stem cells used to grow additional stem cells that match the ones that came from

CAREERS

Cell Biologist in Action

DR. GAIL MARTIN

TITLE Professor, Anatomy, University of California, San Francisco

EDUCATION Ph.D., Molecular Biology, University of California, Berkeley



In 1974 Dr. Gail Martin was working at the University College in London when she made a huge advance. She developed a way to grow stem cells in a petri dish. These fragile cells were hard to work with, so Dr. Martin's breakthrough removed a big obstacle to stem cell research. Seven years later, she made another key discovery while working in her own laboratory at the University of California, San Francisco, in her native United States—how to harvest stem cells from mouse embryos. Her work has helped other scientists develop ways to harvest stem cells from human embryos and explore their use in treating disorders.

Dr. Martin likes to point out that her work shows how small advances in basic biology can pay off years later in unexpected ways. She states that many people focus on cures for specific diseases, not realizing that these cures "may come from basic research in seemingly unrelated areas. What is really going to be important 20 years from now isn't clear."

Read More >>

at HMDScience.com



This researcher is micro-injecting mouse stem cells into fertilized mouse eggs to be used in drug research.

the original embryo. Each source presents its own special set of ethical considerations.

Transplantation into the target area The delivery of stem cells to targeted tissues can be complex, especially if the tissues are deep inside the body. And once delivered, stem cells must "learn" to work with other cells. For instance, inserted cardiac cells must contract in unison with a patient's heart cells.

Prevention of rejection Stem cells may be rejected if a patient's body sees them as foreign. This problem can remain even when certain identifying proteins are removed from the cells' membranes. The development of SCNT technology in humans could help solve this problem so that patients would not have to take drugs to suppress their immune system.

Suppression of tumor formation By their very nature, stem cells remain undifferentiated and continue to divide for long periods of time. When transplanted into an organism, many embryonic stem cells tend to form tumors. This risk must be removed before the cells can be used therapeutically.

Unanswered Questions TEKS 3A, 3D

Stem cell research and therapy do not only involve questions of what we can do. They also involve questions about what we should do, who should benefit, and who should pay.

- Should human embryos be a source of stem cells?
- How should stem cell research be funded?
- How can the benefits of stem cell research best be shared by all people, regardless of income?
- Should insurance cover costly stem cell procedures?

Read More >> at HMDScience.com